

Client: JUNG, KAYE (03182023)

Gender: Female

BATINGA ANIMAL MEDICAL

Patient Name: WOO-YOU

Weight:

CENTER

Species: Canine

Age: 2 Years

Breed: Pomeranian

Doctor: VINCENT RAFAEL SITON

Test	Results	Reference Interval	LOW	NORMAL	HIGH
------	---------	--------------------	-----	--------	------

Catalyst One (March 18, 2023 9:10 AM)

Progesterone 10.8 ng/mL

Progesterone (ng/mL)

< 1.0 Anestrus/Proestrus

1.0-1.9 Pre-LH surge

2.0-3.0 Associated with LH surge, about 24-48 hours prior to the ovulation wave

3.1-4.9 Post-LH surge, pre-ovulation

5.0-12.0 At or near ovulation (progesterone concentration may vary with breed and size)

Physiologic variation in progesterone occurs prior to ovulation. We recommend confirming a continuing rise in progesterone (every 24-48 hours until ovulation is confirmed) when evaluating for timing of breeding.

When measuring progesterone to evaluate for neuter status or for ovarian remnant in dogs (2 weeks-2 months post signs of estrus), a progesterone > 2 ng/mL is consistent with the presence of a functioning corpus luteum and the presence of ovarian tissue. In addition to progesterone concentrations, anti-mullerian hormone, LH concentrations, vaginal cytology, clinical signs, abdominal ultrasound and estrogen concentrations can be used to further evaluate neuter status and for ovarian remnants.